

Paper March 14 1829

Phytochemia,
or
A Dissertation
on *Inland vegetable Chemistry;*
with researches
on the proximate principles of plants;
Submitted, as an inaugural thesis,
to the examination of
the Medical Faculty of the University
of Pennsylvania,
by Columbus C. Corwell of Pennsylvania
March 1829.
South West Corner of Fourth & Spruce



Pensa e mira
Quanto è varia natura, e quai tesori
offre al' uom che la studia e che l'ammira.

Inscribed
to my Receptor,
Dr Samuel Jackson,
As a testimony
of the sincere regard,
and respectful attachment
I have ever entertained for him.



was
was
glo
I was
these
these
the
ance
nities
so on
coming
To co
to se
if th
To ar
their
these
To as
their
I to

These numberless varieties of animals that
wander on the earth, dwell in the wave, or soar
upon the breeze; these stately trees; these delicate
flowers, whose fragrance enchants the sense,
& whose rich tints embellish the surface of our globe;
these burnished metals; these sparkling gems;
these stupendous rocks, at whose crystallization
the lightnings attended, & the waters were
auxiliary: these marvellous combinations, infi-
nitely varied, infinitely modified, these are
so many links in the sublime chain of entities
composing the universe.

To compare them in the range of association,
to study their contrast, & to feel their harmonies,
is the object of the Philosopher.

To arrange them into kingdoms, to determine
their orders, to fix their genera, & to subdivide
them into species, is the duty of the Naturalist.

To analyze their structure, to resolve them into
their principles, to ascertain their properties,
& to isolate their elements is the office of the Chemist.

[Faint, mirrored handwriting, likely bleed-through from the reverse side of the page.]

the
it
y
scu
m
not
we
sub
wor
cap
can
una
In
laia
ham
soph
may
gen
these

Mineral Chemistry has nearly monopolized the attention of investigators, probably because it is more exempt from confusion in its data, & less complex in its products, than its sister sciences; for such elements as analysis can isolate, synthesis can recombine. This is not the case in vegetable Chemistry, nor can we ever hope to form in our laboratories substances developed by the long & incessant work of organization — we may destroy but cannot reproduce — we may analyze but cannot effect a similar assemblage of particles, unaided by the phenomena of vitality.

In the mineral kingdom, substances may be said to combine 2 by 2; in the vegetable, 3 by 3; namely, oxygen, hydrogen, Carbon; & in the animal, where this gradation of complication is at its maximum, 4 by 4 H_2O ; oxygen, nitrogen, hydrogen, Carbon. The chain of affinities connecting these principles is less durable as their complica-

Calio
king
mena
in the
ple
mole
The
be do
object
anism
with
furn
source
of An
of obs
delic
Stim
acid, o
the su
which

lation increased; Hence in the two last mentioned kingdoms are observed the spontaneous phenomena of fermentation in the one, & putrefaction in the other, or the cessation of Life—that principle which, as it were, agglutinates the organic molecules.

The importance of vegetable Chemistry may be duly appreciated, when we reflect that the object of our study is one which confers on Animals the blessing of life, by presenting us with every species of nutriment, & probably furnishing our atmosphere with its exclusive source of oxygen. This harmonious dependence of Animals on vegetables, & vice versa, is worthy of observation. They exhale oxygen, a fluid deleterious to vegetables; we inhale it, as a stimulant to vitality. They inspire Carbonic acid, as an aliment; we expire it, as a poison. We sustain life by consuming substances which they reject as superfluous or diseased;

the
of a
two
above
disting
In
esper
of the
get
ratio
impe
with
the d
corde
"Hogge
into p
"infla
mper
all the
desera

they flourish best in the experimental matter
of animals. This sublime harmony blends the
two kingdoms so intimately together, that the
absence of the one would infer the consequent
destruction of the other.

Indefatigable & successful as have been the
exertions of Chemists in the petetic department
of this science, it is singular that no one has
yet presented us with a judicious or even
rational classification—an object of primary
importance in every branch of study; for
without a definite arrangement we labour in
the dark, & all our inferences are liable to be
considered vague erroneous & empirical. True,
"Vegetable Chemistry" has been arranged by some
into products "Soluble & insoluble"; by others into
"Inflammable, and uncombustible"—these & a
myriad of others, equally philosophical merit
all the popularity they have obtained, & scarce
deserve to be noticed. I have seen one scheme

of a
imita
elegan
coloze
correc

It is
Phyo
their
sion-
trunk
surfac
these
I want
of Rom
in the
by wh
the pic
tound
and the
anethr

* Plum

of arrangement which I intend partially to imitate: It was composed by J. F. Virey, an elegant ^{French} writer, and accomplished pharmacologist; it is a bold & happy attempt at a correct onomatopoeism. bad taste

It is foreign to my design to touch on the Physiological phenomena of vegetables, their phases of increase—their periods of declension—their method of germination—their roots—trunks—branches—their leaves, whose superior surface is exhalant, & whose inferior is absorbent—these researches belong to the botanist: Nor shall I wander with St. Pierre* thro' the poetic maze of Romantic enthusiasm, to admire with him, in the simple flower, the anthera suspended by white fillets poised by double rafters of gold on pillars of ivory; the corolla, an arch of boundless magnitude, adorned with the ruby and the topaz; the glands, superb flagons of amethyst pouring from their gullets of diamond

* Etudes de la Nature

ing
it con
which
pross
The p
furnu
Carbo
Mois
that
it fin
Squid
life, w
In the
sun-p
begec
deligh
(oxid)
of spre
exhale
Navas

ingots of liquid gold. As far however as Chemistry is concerned in the development of some changes which occur in plants, I shall confine it my province to speak.

The principal sources of nutriment are furnished to Vegetables by water, heat, light, Carbonic acid & perhaps earth.

Moisture is a Stimulant so essential to plants that without it they languish & die. This fact is finely illustrated in the case of the onion, Squill &c which perform the functions of life, while suspended in a moist atmosphere. In the burning deserts of Arabia, whose soil, sun parched & sandy, affords no vestige of vegetation, the wanderer's eye is occasionally delighted with the prospect of Isles of verdure (oasis) which appear around the margin of springs. The sunflower is said by Ray to exhale a quart of water in 24 hours; and Navarette in his account of China makes

the
fruit
seeds
vine
refr
the g
decor
the h
way
Heas
natur
polas
Couna
while
only
region
and M
groves
Lapla
winces

Mention of a vine, called the Bejuco, from whose fruit or melon, when punctured, there exude several quarts of cool limpid water. The grape vine & water with of Jamaica afford a clear refreshing water on being cut. Passing thro' the assimilative organs, water suffers decomposition; the oxygen is evolved, while the hydrogen remains to form the base of oils, wax, resins &c.

Heat is also a necessary Stimulant. Vegetable nature is more vigorous in tropical than in polar climates: hence we find that warm countries abound in large trees, and forests; whilst shrubs, underwood, & arbusts are the only kind of vegetables met with in cold regions. Nay the very plant whose size and majesty render it the pride of Turkish groves, when transported to the soil of Lapland degenerated, becomes small, wrinkled, & contracted: because, the rapidity

of day
of the
Sign
on be
lamp
sim
East
after
alime
exper
questio
and in
pound
ion of
rated
it with
yard.
was for
nine po
owned.

of digestion is proportionate to the intensity of the heat.

Light possesses considerable influence on vegetation. The ray emanating from a lamp produces an effect analogous to the ^{the} sun-beam.

Earth. Can the substance of the earth, tho' often regarded as the most indispensable aliment, be considered nutritious? An ingenious experiment of Van Helmont seems to set the question at rest. He dried 200 pounds of earth, and in it planted a willow weighing two pounds. In order to secure it from the admission of other earth, he covered it with a perforated tin hemisphere, carefully watering it with distilled water twice a day, for five years. After the lapse of that period, the willow was found to weigh one hundred and sixty nine pounds, & the earth had lost but two ounces. From this experiment it is certain

that
tal m
just
Chem
the w
of Pot
Plash
eased
not pro
really
develop
assisted
in plan
we heith
our acia
believe
tion? I c

*Lanthorn
of potash
a few veget

that vegetables receive little or no nutrimental matter from the earth. The experiment just cited gives origin to a more important Chemical question: from what source could the willow have derived the many pounds of Potassa which it must have contained? Potash is not volatile it could not have procured it from the atmosphere, and certainly not from the distilled water. Is potassium really elementary, or could it have been developed in the plant by gaseous combination assisted by organic action? Does potash exist in plants during vegetation*? If so, why can we neither detect it by our tests, nor extract it by our acids? If not, have we not every reason to believe that it is literally generated by combustion? I confess myself sceptical enough to

*I do not here speak of that scarce appreciable or doubtful opinion of potash said to be occasionally found in combination with a few vegetable acids.

do
with
of Rea
combu
I mea
Carbo
Puffe
of Pol
enough
in hy
It wa
risk o
have o
degree
combu
The Po
Oreber
prece

doubt whether potash exists ready formed with Charcoal; for how comes it that after extracting from Charcoal all matter capable of Reaction with the acids, we can afterwards convert it partly into potassa by ignition & combustion? I feel perfectly convinced tho' I never performed the experiment that Carbon precipitated from Cyanogen or from Prussic acid, would if burned leave a residue of Potash: and who would be hazardous enough to maintain that the alkali preexisted in hydrocyanic acid?

It would be deemed romancing, in me to risk the hypothesis that charcoal & potassa have the same metallic base, differing only in degree of oxidation; and yet they are both combinations of oxygen with a known base. The Potassium of Davy & the Carbonium of Doberseiner are described with properties precisely analogous, & I am confident, they

are
the ca
partl
Carbo
of Ohio
insolu
Metals
vegeta
my fir
if not
from
Moraea

a. P. P.
bacteria
guishe

b. Guiton
french
prose the

are identic; for during Scheele's process the charcoal was oxidized & converted partly into potash from which the base Carbonium was obtained. The prosecution of this digression from my subject would involve a discussion on the elementarity of Metals; and tho not strictly connected with vegetable Chemistry, I cannot help expressing my firm belief that it is highly probable, if not unquestionably certain, ~~that the Metals~~ from the experiments of Priestley,^a & Guyton Morveau^b as well as from the observations

a. Dr. Priestley, reduced copper to a light combustible charcoal which he had distinguished by the name of Metallick Charcoal

Vide Encyclop: article - Charcoal.

b. Guyton Morveau, in a paper read before the french institute, Floreal 6th Anno 3^d, attempts to prove that the alkalis are combinations of simple gases.

of the
and
each
The ab
the m
of the p
abstra
Had sh
advan
in the
with s
what
apple,
flung
treasure
the h

*Hulloa,
convict
arizon
in the de

of Alloa* that Metals are not simple bodies, and that Chemistry will some day or other teach Men to analyze & to reproduce them. The attempt of the Alchemists to discover the method of making gold, the lapis lazuli, & the philosopher's Stone, was not half so absurd as our philosophers would have it. Had they possessed, like us, the sweeping advantages of analogy & induction, their investigations might have been crowned with success: but accident ^{may} develope what study & research never can. Newton's Apple, Galvani's frog, & a Child's lens, have flung open the doors of the most gorgeous treasury of science that has ever enriched the human intellect.

* Alloa, the Philosophic traveller, declared his conviction that the elementary parts of gold are arisgen from many curious circumstances detailed in the second volume of his voyages.

Yh

I s

Clay

Order

Order

alter

negat

brillia

two of

8. Iod

Genus

Genus

Genus

Genus

Genus

The following is the Scheme of Classification,
I shall adopt throughout this treatise.

Class. Phytochemia from gulos a plant.

Order I. Inland Phytochemia.

Order II. Marine Phytochemia.

It is much to be regretted that chemical
attention has never been directed to marine
vegetables. They certainly present a wide &
brilliant field for philosophical research —
two of our most interesting products, Soda
& Iodine are derived from the sea weed.

Division of order I:

Genus I. Anthracia, from anthrax, Carbon.

Genus II. Hydrogenia.

Genus III. Oxygenia.

Genus IV. Nitrogenia.

Genus V. Hydroxygenia.

Un
pro
me
Sp
con
bel
Solu
den
of Co
recl
tie o
wh
tans
Bra
Con
perh
can
Ligh

Genus I. Anthraxia.

Under this head are comprehended all proximate principles in which Carbon predominates.

Species I

Lignin. The dense fibrous tissue, constituting the parietes of the vegetable vessel, after desiccation & extraction of all Soluble matter by water & alcohol, is denominated Lignin. It contains 52 parts of Carbon in 100, affords to distillation acetic acid combined with an empyreumatic oil (pyroligneous acid) and produces, when treated with nitric acid, artificial tannin, colouring matter & oxalic acid. Braconnot's experiments relative to the convertibility of lignin into sugar are perhaps the most curious & interesting ever recorded in the archives of Chemistry. Lignin, heated with its own weight of caustic

pro
and
Spe
the
it a
com
its c
being
(the
Brid
affo
its d
it ha
By d
arom
Suber
disqu
termes
Spe

potash, affords a substance of easy solubility,
analogous to ulmin.

Species. 2.

Suberin. This term is applied to
the outer bark of the cork tree. From Lignin
it may be distinguished by its greater
combustibility, its specific lightness, &
its compressibility; but chiefly by its
being convertible into a peculiar acid substance,
(the suberic) by reaction with nitric acid.

Boiled in alcohol, & evaporated suberin
affords a crystalline principle, called by
its discoverer, Cerin — for distinction sake,
it has since been entitled Suber-Cerin.

By distillation, I procured from cork an
aromatic acid which may be called pyro-
suberic with the same propriety as the
disguised acetic acid of Lignin has been
termed pyroligneous.

Species. 3.

G
pr
exa
Sub
Sp

pro
Cous
also
Nitin
Map

prin
wall
of Lin
Nitin
separ
sema
had

Gossypin. Cotton is certainly a peculiar principle. It has been very trivially examined. With nitric acid it affords an acid substance resembling the Induric.

Species 4

Mucin. The tissue of the Mushroom possesses properties which entitle it to a separate consideration. It is inelastic, insoluble in water, alcohol, or ether, soluble in the alkalies & in nitric acid, which converts it into a ceruminoid mass.

Species 5.

Tannin. The vegetable astringent principle may be obtained by pouring lime water on a strong solution of tan. The tannate of lime is formed & may be decomposed by nitric acid, which uniting with the lime, separates the tannin. After filtration there remains a black pulsatent substance which has been described as pure tan (Krey)

Acetone
be it
it from
of Am
tan is
with y
soluble
of tan
which
Artific
nitric a
vegeta
Speed

Greek o
garden a
the long
form a

four

According to Boast & Lagrange, Tannin may be isolated in all its purity, by precipitating it from an infusion of Galls by Carbonate of Ammonia. The principal peculiarity of tan is that of forming an insoluble compound with Gelatine. It is a brownish black powder, soluble in water. There is, however, a variety of tan existing in Cinchona & other tonic barks which is soluble in alcohol alone.

Artificial tannin is formed by reaction of nitric acid with Charcoal lignin & several vegetable substances.

Species 6

Chlorophyllin. This term is of Greek origin being derived from *Chloros* green, & *Phyllon* a leaf. The green colouring matter of the leaf, erroneously represented as a resin, forms a peculiar principle abounding in

* Journal de Physiologie par Magendie - Janvier 1821.

Hy
Sub
also
also
Sub
he dis
it is
is so
red by
altera
yellow
nitric
either
Spec
errone
but ca
the m
a qu
termed

Hydrogen. It may be obtained by digesting buds or young leaves in highly rectified alcohol; after a slow evaporation of the alcoholic solution, a splendid green substance remains. Thus obtained it may be distinguished by the following properties: it is soluble in alcohol oil & ether, insoluble in water & burns like a resin; it is discolored by Chlorine & Iodine, is dissolved without alteration by sulphuric acid, strikes a bright yellow hue with Hydrochloric, & a grey with nitric acid, unattended by the production either of Mucic or oxalic acids.

Species 7.

Gumresin. This term is obviously erroneous, being not only vague & gratuitous, but calculated to ~~mislead the~~ impress on the mind the idea of a compound formed of a gum and a resin. Sugar might be termed a gum resin as properly as gamboge.

These
products
of Red
of 10
not a
left
All
are
from
into
soluble
Conce
tanno
under
by ind
A. H. A.
A. H. A.

There exists not in Nature, nor can there be produced by art, any immediate combination of Resin & Mucilage without the interposition of some chemical reagent. Why then will not the Chemists furnish us with a title less calculated to mislead?

All substances classed under this head are obtained by spontaneous exudation ~~or~~ from natural fissures or incisions made into ~~the~~ plants which afford them. They are soluble in water oil & alcohol, & may be converted by the stronger acids into artificial tannin & charcoal. They appear at first under the form of a milky juice, & subsequently inspissate; such for example are aloes, Asafetida, Gamboge, Opium, Galbanum, Lactarium, Scammony, &c.

St
under
pred
gran
Sp

a. S
rege
enton
at ip
with
huga
Bred
Surpr
Comm
Lee-hu
hexago

parman

Genus II

Hydrogenia.

All proximate principles, comprehended under this genus are inflammable, from the predominance of Hydrogen. Their specific gravity is less than that of water.

Species 1

Ceruminous matter.

a. Beeswax. Whether beeswax be a vegetable secretion or the product of entomic elaboration, is a point yet remaining at issue & it is highly amusing to observe with what animated hostility this nugatory contention has been maintained. Beeswax is found on the superior glossy surface of leaves, & on the anthers of flowers; commonly, however it is obtained from the bee-hives in which it is found disposed in hexagonal cells. Sparman^d suspects, without

Sparman's voyage to the Cape.

Ma
Corrie
Du K
Call
a wh
Kase
in the
them
"Quilch
A St
does
is the
Kessu
in wo
fusih
the a
Hear
Spher
boilin
had O.
* Side Sub

much reason, that it is deposited on the
berries of the *Myrica Carifora* by insects; &
Du Halde^{*} alleges that a tree in China
called *tong sing* is absolutely coated with
a white wax formed there by insects. But
travellers are naturally chartered to deal
in the mythic & the marvellous, and to
them as well as to painters & to poets,
"Quilbet auctendi semper quitaqua potestas!"

A sterling Chemical evidence that wax
does not belong to the animal kingdom
is the absence of Nitrogen.

Bee-wax is straw coloured, insipid, insoluble
in water, soluble in the fixed oils, inflammable,
fusible at 150, rendered saponaceous by
the alkalies, ^{unaffected by} ~~unaffected by~~ ^{unaffected by} the acids, and
bleached by exposure to Chlorine & to atmos-
pheric air. It is partially soluble in
boiling alcohol, water, & ether. Dr. John
has contrived the term *Cerin*, to denote

* vide Du Halde's China.

that
the ap
& the
residue
it is so

T
few
a view
the act
Imme
portion
into so
on the
whole
Semi
Treated
Chronic
develop
Boile
and cor

that portion of wax which dissolved in the aforementioned menstruum, when hot; & the term Myrcerin, to designate the residue which under all circumstances is insoluble.

The following is the result of a few experiments which I performed with a view to ascertain whether leucowax resisted the action of our stronger chemical reagents. Immersed in boiling nitric acid, a minute portion of it, probably, the Cerin, entered into solution, the rest remained fluid on the surface, and on refrigeration the whole of it became concrete, white, and semidiaphanous.

Treated in the same manner with hydrochloric acid, similar phenomena were developed.

Boiled with sulphuric acid, its tissue and cruminous character were destroyed.

it a
Sapo.
that of
being
About
with on
Comb
than w
lustre, a
into a Sap
Sapo
to a Sing
in all its
it from S
was will
a gelatin
bath & a
instant
limpid, &
boiling do

it acquired a greenish black colour, a saponaceous greasy feel, & a taste like that of an acid soap, its inflammability being materially impaired.

About twelve parts of wax were heated to fusion with one of Iodine in a Florence flask; a peculiar combination resulted. This compound was heavier than water, more brittle than wax, assumed a metallic lustre, and displayed a rich dark purple hue. Formed into a taper with cotton wick, it burned with great splendour.

Sapocerin. I have presumed to attach this name to a singular compound of acid and alkali analogous in all its chemical relations to beeswax. I obtained it from soap by the following process: fine white soap was triturated with water, until the whole acquired a gelatinous consistency. Being heated in the sand bath & a few drops of nitric acid poured upon it it instantaneously separated into two portions, the limpid, the other opacuminous and insoluble. After boiling down for some time, I separated a yellow

made
redon
every
saffie
Crimo
in eye
Satisf
Subbl
with
Lapoe
by the
water
about
Adip
room, a
alcohol
Spec
by
by Col
propert

matter separated on cooling. From the striking resemblance it bore to beeswax, I repeated on it every experiment I had performed on the subject of wax, in order to ascertain whether it would coincide with wax in chemical analogy, as well as in external appearance. My results were decidedly satisfactory; & justified me in believing, that this substance was, with trifling modification, endowed with every property and character of beeswax.

Sapoeerin is of a reddish yellow colour, ^{insusceptible of} unaffected by the acids, slightly unctuous, insoluble in water alcohol or ether; its specific gravity being about .96, & its point of fusion below that of beeswax.

Adipocerin, discovered by Braconnot in the musk room, is solid, white, & entirely soluble in boiling alcohol, from which it separated on refrigeration.

Species 2.

Vixed oils. These oils are commonly obtained by cold pressure & may be known by the following properties: They are viscid, insipid, immiscible

with
hot, for
photo
of a sum
conceal
8 resin,
which
concrete
Two
have be
X pres
During
produces
lime oil

The deom
or vegetal
is allage
Lancidic
from a s
is absorb

with water, lighter than water, heavier than alcohol, form soaps with alkalis, dissolve sulphur phosphorus caoutchouc & the resins, are incapable of assuming the aeriform state without decomposition, convertible by weak nitric acid into a kind of wax & resin, and may be decomposed by chlorine, which uniting with their hydrogen renders them concrete.

Two immediate principles, elain & Stearin, have been separated from these oils by congelation & pressure between folds of ~~the~~ spongy paper. During the process of saponification, Stearin produces Margaric, and elain oleic acids.

Olive oil consists of Stearin 28
Elain 72

100

The decomposition of the sweet principle of Scheele, or vegetable Muriatic, which all these oils contain, is alleged by some to be the cause of their rancidity. Ist Have supposed rancidity to result from a slow combustion during which oxygen is absorbed.

The
the
from
ture
attrib
lation
Hered
Some
oleage
with
dry
Tegla
Stage
Chen
differ
compos
at atm
Consider

The oxidation of the oil may be accelerated by the decomposition of the mucus.

From the necessary presence of a certain temperature, & the rapid generation of acetic acid, I should attribute the phenomenon to the process of fermentation, which indeed is but another term for Fermentation.

Some oils, by exposure to the atmosphere, lose their oleaginous character, inspissate, & become covered with a pellicle; these are called the siccatives or drying oils.

Vegetable Butters might be mentioned in this stage of the discourse, but as they coincide no striking ~~characteristic~~ peculiarities, & as I conceive they differ from fixed oils merely by being principally composed of Stearin which renders them concrete at atmospheric temperatures, I shall waive the consideration of them.

Sp

before
by app
such
that
As they
from a
Hapth
dormant
instrus
a lion
but ha
of the
They a
normal
disting
their co
their or

Species 3.

Etherial oils.

There is a class of oils, never before noticed, which can neither be obtained by expression nor distillation; but by digesting such substances as afford them in Ether, & leaving that Menstruum to spontaneous evaporation.

As they differ in appearance & chemical relation from all other immediate oils, resembling rather Naphtha or petroleum, I have thought proper to denominate them, from the fluid thro' whose instrumentality they were obtained, Etherial oils - a term formerly synonymous with volatile oils but happily for my purpose that application of the term is obsolescent.

They are yielded by all articles in which the aromatic quality predominates, and may be distinguished by the following properties: Their consistency is bituminous, their ^{colour} ~~odour~~ varying, their odour penetrating, & their taste intolerably

with
with
for
for
or
Eth.
St. X
I Ma
hally
cation
Eth.
of Ph
perce
not to
action
Eth.
thous
Himac
Eth.

warm & acrimonious! They afford to distillation an impure acetic or empyreumatic acid in combination with a volatile, or more properly a pyro-etherial, oil, far less powerful than themselves; a considerable portion of their aromatic principle being decomposed, or obfuscated by assuming the gaseous form.

Etherial oil of Capsicum is perhaps the warmest & most penetrating substance known in Nature. I make no doubt but it will be applied internally to disease, or to purposes of endermic medication with the most decided advantages.

Eth: oil of Pepper, obtained by Mr. G. W. Carpenter of Philadelphia, concentrated all the acrimony and penetrating heat of Black pepper & to this principle, not to the piperin, is to be referred the stimulant action of the substance producing it.

Eth: oil of Cloves communicated to the smell an odour highly fragrant, & to the taste an agreeably stimulating warmth.

Eth: oil of Mustard is composed chiefly of Stearin

With a
Caraway
and the
herb
sensitive

the plant
Lycopericon
Most of
substance

Spe

Most of
in the
substance
obtained
herb
a warm
temper
Aroma
The 11th.

With regard to the Eth: oils of Allspice, Ginger,
Caraway, Calamus aromaticus, Nutmeg, Cubebs,
and others which I obtained, it is only
necessary to state that they embody all the
sensible & medicinal properties of their respec-
tive plants.

Synchronously with the development of
most of these oils, a neutral crystalline
substance appears on the sides of the vessel.

Species 4.

Volatile or Pyroethereal oils.

Most of the volatile oils are obtained by distilling
in the Italian recipient, with water such
substances as produce them; while a few are
obtained by expression, as oil of lemon, orange, and
bergamot. They possess a penetrating odour &
a warm aromatic taste, are volatilised at common
temperatures, and pass into resins, & lose their
aroma by oxygenation.

The Vol: oils are divided by Virey into 3 orders:

1st the

Best of

2nd the

Most of

3rd the

for the

in the

These

bottoms

by sugar

its

applies

inflame

a spongy

mark

acid gas

smaller

had been

- 1^o the light oils, as oil of thyme, rosemary, orange &c
Most of which deposit crystals of Camphor;
- 2^o the heavy, as oil of Cinnamon, Cloves, Mace &c
Most of which deposit crystals of Benzoin Resin;
- 3^o the concrete, as butter of roses, anniseed, fennel &c
for the distillation of which more heat is required
in the refrigerant vessel.

These oils are all empyreumatic & acid: they render
bitterness & may be rendered miscible with water
by sugar or mucilage. They precipitate gold in
its metallic state from its nitro-muriatic solution
deposit Camphor, Sulphur & phosphorus,
inflame by reaction with nitrous acid & leave
a spongy cellulated resin, called the philosophic
mushroom. When a current of Hydrochloric
acid gas is passed thro' a volatile oil, a crystalline
matter analogous to Camphor, falls. This deposit
has been called artificial Camphor.

Spo

solu

fract

such

from

in wa

secta

deco

holic

made

prina

artifi

Roma

extrom

tain:

soluble

and so

bitter

& near

Species 5.

Resins may be regarded as concrete volatile oils. In a state of purity, they are friable, vitreous, inflammable, insipid, & inodorous. Such of them as possess sapidity or odour derive it from an essential oil or benzoic acid. They are insoluble in water, soluble in oil, alcohol, ether, acids, & alkalis. Acetic & Hydrochloric acids dissolve them without decomposition, Sulphuric acid destroys them, & Nitric acid converts them into a bitter yellow matter, entitled, in honor of its discoverer, the "Bitter principle of Welcher"—a portion of Mucic acid & artificial tannin are simultaneously produced. Bonastre, who analysed the natural resins with extreme accuracy & minuteness, found them to contain: 1° a resin, properly so called, translucent and soluble in cold alcohol; 2° an acid, sometimes acetic and occasionally succinic; 3° an essential oil; 4° a bitter matter; 5° a subresin, opaque, crystallizable, & nearly insoluble in boiling alcohol. This last

appear
are to
of Guinea
become
of powder
renders
or by Xim
of Porcelain
of a paper
sent by
Colours w
The letter
clater in
extracts,
Substan
principle
The torer
fluid by
by distill
with ben

appear to be to resins what elain & Stearin are to fixed oil.

a Guaiacum is partially soluble in water, & becomes successively green, blue, & brown by exposure to atmospheric oxygen. It may be rendered blue by torrefaction with gum arabic, or by Ximome.

b Seruline. This term is applied to the resinous portion of asphaltida gum ammoniac &c. It becomes rufescent by exposure to the atmosphere, & strikes a rose colour with silver.

c The bitter resins, so abundantly found in Colocynth, elaterium, Scammony, & most of the panchymagogic extracts, are tanno-resinoids or combinations of substances resembling tannin with a bitter drastic principle.

d The terebinthinates are resins rendered semi-fluid by an essential oil, which may be separated by distillation. The ballams are resins impregnated with benzoic acid, which may be disengaged by

Sabine
Spe

be reg
and bo
and sh
like the
indol
Camp
diaph
proper
& putty
jenu
distill
group
Camp

This ge
prin

Sublimation.

Species 6

Camphor. This singular substance may be regarded as intermediate between resins and volatile oils: like the former, it is light solid and inflammable soluble in alcohol oil and ether like the latter, it is volatile, aromatic, & nearly insoluble in water.

Camphor is concrete, crystalline, white, semi-transparent, volatile, & combustible. It has the property of arresting the progress of fermentation & putrefaction in vegetable bodies. Its principal peculiarity is that of forming by repeated distillation with nitric acid an acid substance, grouped in plumose crystals entitled the Camphoric.

Genus III

Nitrogenia.

This genus comprehends all immediate principles containing Nitrogen.

In
shall
lack
ation
Kouta
J

posterior
Starch
paste
It was
K. May
is a fl
in mod
Luscep
After a
and w
acid, Ca
pyrogo
superpe

For the sake of a definite nomenclature, I do
shall annex the ~~the~~ termination al or ia to
such principles as are alkaline, and the termin-
ation in to such crystalline products as are
neutral.

Species 1

Gluten. This name is given to that
portion of farina which remains after all the
starch is washed away, by forming it into a
paste & decocting repeatedly with cold water.
It was first observed by Beccaria & Kessel Meyer,
& may be known by these properties viz: it
is a flabby, tenacious, grey substance soluble
in most of the acids & caustic alkalis, and
susceptible of acetous and putrid fermentation.
After desiccation it becomes hard & vitreous
and when heated produces ammonia, phosphoric
acid, cacic acid, sulphuretted hydrogen, and a
pyrogenic oil. With alkalis it forms an
imperfect soap, and affords with alcohol

who
saw
also
with
of his
sphere
diffic
wine
yellow
little
the
prompt
thems
of turn
pursu
& wro
into w
solall
Spee

which dissolved a portion of it, a fine yellow
varnish. The portion of Glutin Soluble in
alcohol has received from Taddai the appella-
tion of Gladine; the insoluble portion, that
of Limone. The latter is found in little
spherules which adhere to each other with
difficulty, and exhale the odour of putrifying
urine. Gladine when dry, is brittle
yellowish & translucent. It is, I find, subdivi-
sible into two portions and were it worth
the expenditure of time ink or paper to attach
pompous academical names to products in
themselves so nugatory (the most effectual method
of turning Chemical Science into ridicule) I might,
pursuant to the example of the Taddaian school,
& with equal propriety subdivide his gladine
into two portions — the one, as a matter of course,
soluble in boiling alcohol; the other, insoluble.

Species 2

a Albumen. This substance bears a

very
It is f
milky
in Cabb
several
destructive
princip
happy
had no
into the
b. Cellar
is incorp
its solu
it form
c. p. m.
Coincide
d. Catatē
parent,
He in
gluten, &

very intimate resemblance to animal albumen. It is found in a few vegetable juices which are milky, and coagulable, by heat or by acids: as in Cabbage, horse-radish, gonio, poppy, lettuce, & several of the Cruciform plants. Submitted to destructive distillation, it affords azotified principles. It is partially soluble in alcohol. Happy for the student of Chemistry, Taddai has not experimented on the soluble and insoluble portions.

b. Gelatine, found in the pith of the date, is incoagulable by heat, and precipitable from its solutions by an infusion of galls, with which it forms an insoluble compound.

c. Myzome, observed by Tanguelin in mushrooms, coincides in all its properties with animal myzome.

d. Caoutchouc is tenacious, elastic, and semitransparent, insoluble in water and alcohol, soluble in Ether, oil & in Nitric acid. It resembles gluten, & exhales Azote and ammonia when heated.

Sp

the
a. 2

pos
disco
ately
the
the
is be
by h
exce
corate
rema
alco
depos
b. Cin
differ
may

Species 3.

Proximate principles which are
alkaline & Crystallizable.

a. Quinia.

The alkali, to which the tonic
powers of quinquina are referable, was
discovered by St-Gomez, and afterwards accurately
examined by Pelletier & Caventou.

The following is M. Pelletier's process for
obtaining it: the alcoholic extract of barks
is treated with hot water rendered acidulous
by hydrochloric acid, and boiled down with
except of Magnesia; after filtering, and decan-
tating with cold water, the matter which
remains on the filter, digesting it in strong
alcohol, and evaporating, pure quinia is
deposited.

b. Cinchonia, as it exists in the pale bark,
differs little from the preceding, and
may be developed by a similar process-

c. 200

detest
in wh
princi
a piece
It wa
powde
in two
brack
for som
and th
solu
On rep
precip
By ex
with
treatin
fili
evapo

c. Quassa.

This new vegetable alkali was detected by me in the Quassia amara, in which it is entailed with a variety of principles, as tannin, colouring matter, a peculiar new acid (~~the~~ quassic) and resin.

It was thus isolated = One pound of powdered quassia was digested a few days in two quarts of water, with about four drachms of Sulphuric acid: after boiling for some time it was filtered while hot, and the evaporation continued, until the solution became perfectly concentrated.

On refrigeration the Sulphate of Quassa precipitated in snow white acicular flocks. By ~~boiling~~ boiling the sulphate in water with excess of Magnesia nearly to dryness, treating the residuum with hot alcohol, filtering to get rid of the Epsom salt, & slowly evaporating the alcoholic solution, I obtained

just
In

energy
which
the Pa
conde
very
and
tuted
salty
least
form

La
He, a
person
quanta
pharm
cum lo
taken,
of gay

pure Quassa.

Quassa concentrates all the medicinal energy and bitterness of quassia. It is white and eminently alkaline, precipitates the metallic oxides, neutralizes the acids, and renders the tincture of alkannet blue: it is very soluble in hot alcohol, less so in cold, and nearly insoluble in water. It constitutes the basis of a class of intensely bitter salts. For so, Quassa is uncrystallizable at least I could never procure it in a crystalline form.

Sulphate of Quassa is easily crystallizable, and grows in acicular prisms. It is a permanent salt, and remains solid in any quantity of cold water. In the common pharmaceutical preparation "infusum quassiae cum sulphate zinci," a double decomposition takes place; the Quassate of zinc, and Sulphate of quassa being formed.

The de
guaratio
Maele
Acclac
fibred.
Opalac
d. S.

Similar
of Duak
Wriston
root, a
powder.
It is d
the sulph
the accl
if not c
product.
The Sulph
the vid

Hydrochlorate of Quassa forms starry configurations, and is more soluble than the preceding.

Nitrate of Quassa is uncrystallizable.

Acetate of Quassa forms in silky plumose fibres.

Oxalate of Quassa aggregated in fine bundles.

d. Serpentina.

By pursuing a plan similar to that indicated for the development of Quassa, I succeeded in isolating, from the *Aristolochia Serpentina* or Virginia Snake root, a new alkali to which the tonic powers of the Virginia root are attributable. It is better in every instance first to procure the sulphate, or what may be more eligible the acetate, Sulphuric acid being liable, if not cautiously employed, to derange the product.

The Sulphate may be disentangled from the viscid extractive matter which obstinately

adher
presen
the last
Magne
led in a
stallio
it appe
deliques
smell a
renders
Sulph
long qua
cold wa
Hydroc
fired more
Nitre of
to speak
large ad was
at least I can
the immense

adhered to it by repeated washing and
pressure between folds of bibulous paper.
This salt being boiled in water with excess of
magnesia, evaporated to dryness and diges-
ted in alcohol, pure *Serpentaria* falls, after
ebullition and filtration. Thus obtained,
it appears in white and exquisitely delicate
deliquescent crystals, which possess a fragrant
smell and a slightly bitter taste, and
render the Symp of violets ~~blue~~ green.

Sulphate of *Serpentaria* crystallizes in
long quadrangular prisms. It is insoluble in
cold water, but soluble in hot water and alcohol.

Hydrochlorate of *Serpentaria* forms brilliant plumbe-
fied, more soluble than the sulphate.

Nitrate of *Serpentaria* is not susceptible of crystallization,
or to speak more justly, nitric acid decomposes *Serpentaria*,
as well as most of the other vegetable alkalies—
at least I could never regain them after having
once immersed them in that acid, however dilute.

Sept
a piece
I have
procured
intelligence
in the C
e. Colo

re from
with a n
of the act
may be
for the
appears
North Sa
and we
deliques
Combined
fine nee

Serpentaria naturally exists in union with a peculiar aromatic acid (the serpentaria) which I have not examined, owing to my inability to procure it in sufficient quantity for purposes of investigation. The serpentaria of lime is found in the Caput mortuum with extractive matters.

c. Columbia,

Another new alkali extracted by me from the Colomba root exists in combination with a new acid (the columbic) tannin, resin, extractive gum, and mucus. From all these it may be disengaged by the process employed for the isolation of Quappa. It is white and appears in the form of shining silky needles. With sulphuric acid, it forms minute prisms; and with hydrochloric, starry points, somewhat deliquescent. Nitric acid destroys its alkalinity. Combined with acetic acid, it forms very fine needles resembling aspidinthes.

f. G.

rated

quasi

differe

ding

and

propo

by the

the Ch

the m

formu

deal of

Studen

fatigue

Gente

Chasa

which

in 1m

which

f. Gentia

A new organic alkali which I separated from Gentian, in which it exists in great quantity. The method of obtaining it does not differ from that employed for procuring the preceding alkalis. The process is exceedingly simple, and I feel thoroughly convinced that all other proximate alkaline principles might be ^{obtained} ~~procured~~ by the same simple method. This would save the Chemists, who seem to vie with each other in the intricacy, prolixity and unintelligibility of their formulae for obtaining these principles, an enormous deal of trouble, and exonerate the mind of the Student from much perplexity, and unprofitable fatigue.

Gentia embodies with little variation, all the characters of Columbia. The Sulphate of Gentia, which like the Sulphate of Columbia appeared in small needleform crystals, and the Muriate, which like that of Columbia appeared in

ad
g. G

colon
Muc

a how
small

Sulph
part

all po
it de

h. St

Wade

disce

his me

have b

publ

that g

on the

astroid points, were the only Salts I examined.

g. *Gallia*.

Besides gallic acid, ellagic acid, tannin, colouring matter, a green mossy substance, Mucus, and extractive, there exists in the gallnut a new vegetable alkali which I obtained in small quantity by Sulphuric acid.

Sulphate of *Gallia* is insoluble in cold water, partially soluble in hot, and dissolved in all proportions by ⁱⁿboiling alcohol, from which it descends in fine fasciculi on refrigeration.

h. *Stramonium*.

This alkali has been mentioned under the appellation of *Daturia*, and its discovery ascribed to Brande; but as neither his method of procuring it nor its properties have been described in any of our Cisatlantic publications, I am warranted in believing that I present the first series of investigations on the subject of an article which will, one

to
to
The
dual
were
the p
With
be op
from
enter
this
be to
Supp
in wo
redig
redig
shall
Stra
which
be pre

day or other, prove an invaluable accession
to the Catalogue of remedies.

The subjoined is the process I employed for its
development. The bruised seeds of *Stramonium*
were washed in cold water and alcohol for
the purpose of removing extraneous matter.
Without this preliminary precaution it will
be extremely difficult to separate the alkali
from the mass of other substances which
enter so abundantly into the composition of
this seed. After repeated elutriation, it may
be treated with dilute sulphuric acid, and the
sulphate obtained. This salt being boiled
in water nearly to dryness with excess of mag-
nesia & boiling alcohol being poured on the
residue, which must be filtered during
ebullition pure *Stramonium* falls on cooling.

Stramonium crystallizes in acicular flocks,
which when minutely examined appear to
be prisms with trapezoid bases.

It is
but a
Sulph
latica
Hypoc
of some
Acetate
The ac
neutral
and is
Stannic
i. Mac

dance
in comb
princip
The Sul
examina
It is ins

It is insoluble in cold water and alcohol
but very soluble in hot alcohol.

Sulphate of Stramonium crystallizes in quadri-
lateral prisms.

Hydrochlorate of Stramonium assumed the form
of small cubes.

Acetate of Stramonium groups in delicate needles.

The acid by which Stramonium is naturally
neutralized exhales the odour of ripe apples,
and is perhaps malic acid.

Stramonium is an energetic narcotic poison.

i. Mahogonia.

This alkali exists in great abun-
dant in the bark of the Sweetgum Mahogoni
in combination with a splendid colouring
principle.

The sulphate, the only preparation I have
examined, crystallizes in long prismatic needles.
It is insoluble in cold, but soluble in hot water.

k. 2

(Que

gab

colo

relos

back

Pha

on a

Quar

that

the s

with

with

l. s

great

febrile

ly, also

of the

k. Quercia,

was obtained by me from Oak Bark, (*Quercus robur*) in which it is united with gallic acid, inert substance, and a peculiar colouring matter (*Quercin*) hereafter to be described. The invigorating properties of Oak bark, gratuitously attributed to tannin in our Pharmacopoeia, are unquestionably dependent on this alkali. I presume from analogy that Quercia is, per se, crystallisable, and feel confident that it will rank among our choicest tonics; the wood which affords it being procured, without expence, and the alkali obtained without difficulty.

l. Angusturia.

This alkaline base I obtained in great quantity from Angustura bark (*Cusparia febrifuga*). The bark furnishing it had been confidently alleged by the most respectable practitioners of the West Indies to possess a decided superiority

1862

Thidg

Sagu

will p

Sagu

deduct

in Mo

Sals

m. Croa

of Sag

which

acid.

Sals

of its

refers

suspec

Suppor

n. Cro

over Cinchona, in the treatment of intermittent fevers.
This granted, we need not hesitate in assuming, that
Angustura, if not more serviceable than Quina,
will at least prove a substitute of equivalent value.
Angustura, to which I have given but a very
desultory & superficial examination, crystallized
in minute deliquescent fibrils.

Sulphate of Angustura aggregates in Stars.
m. Cordia.

This alkali was separated in the state
of Sulphate from the wild-cherry-tee bark in
which it probably exists combined with hydrocyanic
acid. The only combination I examined was the
Sulphate. This however sufficed to convince me
of its alkalinity, and to this principle may be
referred its medical activity & not as Murray
suspects, to the prussic acid which is merely
supposed to exist in the bark.

n. Crotonia.

The alkali, in which reside the fragrance

and
Case
by the
pres
bale
appe
each
s. Ca

alka
Lik
in de
p. Ca

by the
insti
exist
to pipe
line
in con

and tonic energy of the Croton eleutheria or
Cascarilla bark, was detected by me, and developed
by the process so often alluded to. After being
pressed in soaking paper dissolved in boiling
water, and recrystallized, the Sulphate of Crotonia
appeared in long acuminate prisms, interlacing
each other.

o. Canellia.

From the Canella alba I extracted an
alkaline base by means of Sulphuric acid.
Like Sulphate of Crotonia, it presented itself
in delicate interwoven fibres.

p. Capsica,

A new vegetable Alkali discovered
by Mr G. W. Carpenter. After numerous experiments
instituted for the purpose of ascertaining the
existence of a principle in red pepper, analogous
to piperin, he succeeded in developing a crystal-
line substance which, he informs me, is Capsicin
in combination with a peculiar acid.

of Br

by

More

ephi

in g

r. Mo

disco

st m

infel

fellie

id to

laked

in py

s. Br

Cases

and

In the

9. Cornia.

An alkali, yielded in small quantity by the Cornus Florida, and discovered by Dr. L. G. Morton, is reported to have been advantageously exhibited as a substitute for Quinia. It crystallizes in quadrangular prisms.

10. Morphia.

The alkaline anodyne principle was discovered by Serturner in opium, from which it may be isolated by boiling an aqueous ~~infused~~ solution of opium with magnesia, and filtering: the matter remaining on the filter is to be treated with boiling alcohol, which takes up the alkali, and deposits it on cooling in pyramids and four-sided prisms.

11. Brucia.

Was first observed by Pelletier and Caventou in the bark of the Bonia Antidysenterica and in the Myrica and bean of St. Ignace. In the two last, it is found in company with

W
by
sion
in
bad
1

com
with
shall
isola
cry
It is a
a

from
and
v. A

the a

Strychnia, from which it may be distinguished by its being more easily soluble, and less deleterious to the animal economy. It crystallizes in oblique prisms with parallelogrammatic base.

i. *Strychnia*.

The alkali which exists in the *Impatiens* and bean of *St Ignace*, in combination with a crystallizable acid (the *igaduric*) may be obtained by any of the processes employed for the isolation of *Prophria*. It is found in prismatic crystals terminated by quadrangular pyramids. It is a horrible and intensely bitter poison.

ii. *Picrotoxia*

was first separated by Boullay from the *Cocculus Indicus*. It crystallizes in prisms and may be converted by nitric into oxalic acid.

x. *Atropia*.

It has been a point at issue, whether the active narcotic principle of *Belladonna*

be
of the
The p
in S
and p
Both
time
w. S
task
and
x.
solar
space
line
gation
of bird
Hyo
alkal

be alkaline. It is decidedly so — I procured two of its salts the Sulphate, and the Muriate. The former is soluble in water, and crystallizes in slender prisms; the latter is nearly insoluble and forms brilliant silky needles.

Both these salts, as well as Rhopia, are destructive of animal life.

v. *Daphnia*,

detected by Vauquelin in the bark of the *Daphne alpina*, saturates the acids, and forms a class of Neutral Salts.

x. *Solanum*,

Obtained by Desfosses from the *Solanum dulcamara*, in the form of a white opaque powder, possesses but feeble claims to alkalinity: its salts are all insusceptible of crystallization, and the change effected by it on Litm^{us} of violet is scarcely appreciable.

y. *Hyoscyama*, *Cicuta* & *Acornita*, the supposed alkaline bases of black hellebore, hemlock, and

Part
Every
by
low
able

Acc

exp
The
and
prop
form
by a
add
conce
add
falls

Monk's hood, were reported some time ago, in the European periodicals, to have been discovered by Baernde. They have not however stood the test of rigid scrutiny — no chemist has since been able to procure cicuta.

Species 4.

Poysimate principles neither acid nor alkaline.

Piperin.

This term is given to a neutral crystalline matter existing in black pepper. Mr. G. W. Carpenter, who published a lucid and elaborate memoir on the preparation and properties of this article, employs the following formula for its isolation. It may be obtained by digesting pepper in alcohol, evaporating, adding very dilute hydrochloric acid, and concentrating the solution by evaporation: on adding potash to neutralize the acid, piperin falls. It has been extensively employed

in a
tonic
and
poet
Shor

Sona
tally
scale
attra
The p
colo

inert
Mr. M
the p
being
book

in intermittent fever, & is said to be an invaluable tonic. Per se, however Piperin is totally inert and is indebted for its tonic property to a portion of ethereal oil, almost inseparably adherent to it.

Cathartin.

The purgative principle of Senna is yellow, "transcaborde", and uncrystallizable, soluble in any proportion in water and alcohol, insoluble in ether, and attracts humidity from the atmosphere. The persulphate of iron strikes a fine brown colour with its aqueous solution. (Lassaigues)

Rhubarbacin.

A barbarous term for an inert powder, found in Rhubarb, which Mr. Hanl, its discoverer, very prudently cautions the public to use but little of it; two grains being sufficient for a dose. Mr^r Carpenter took twenty grains of it with no more effect

Cha
Jal

Such
Ehu

few p
Cotto

Wap
N

so de
fabr

which

Nich

if pop

choy

whose

indica

See Bay

than if he had swallowed as much Hæsch's
Jalapin.

The discovery of this substance forms
such another epoch in chemical science as
Ehubarbarin. Tho' the gazettes eulogized it, and a
few practitioners extolled it as the very flower of
Cathartics, it has been found to be Sulphate of
Ammonia.

Nicotin.

It is inconceivable how a Chemist
so deservedly illustrious as Mr Vauquelin, could
fabricate a process so palpably absurd as that
which he has contrived for the developement of
Nicotin; and what appears still more anomalous,
if possible, is that Chemical compilers should
thoughtlessly adopt and republish an operation
whose glaring inconsistencies a child might
indicate.

* See Carpenter on Rhubarb. Chapman's journal for Feb. 1828.

aff
of so
whic
holic
In a
and
whic
fil, it
than
time

C
C
N
S
M
P
Y
w
with

The following is a list of essential principles, some of which I obtained in crystals and all of which may be easily obtained, by the process noticed for the development of the ethereal oils.

In a state of purity, they are nearly insipid; and as all the remediate energy of substances which furnish them depends on an ethereal oil, it will be superfluous to mention more than their names, and indicate at the same time the plants affording them.

Capsicin	from	Capsicum.
Cubebin		Cubeb.
Calamin		Calamus aromaticus
Caryophyllatin		Cloves.
Zingiberin		Ginger.
Sinapin		Mustard.
Myristicin		Mace.
Piperin		Pepper.

These substances invariably exist in combination with Eth: oils but are never found in company

with
Loring
2nd

My a
to bo
tast
challe
with
guar
Quere
in ho
St. Sta
rior in
the id
object
dys.

Nar

disco

with Essential oils being destroyed by the distilling heat.

Quercin.

This substance I obtained, during my attempt to separate Quercia from oak bark. On boiling the rasped bark in water, rendered tart by Sulphuric acid, and filtering during ebullition, a yellow powder, (Quercin) mixed with the alkali of oak, precipitated in great quantity on cooling.

Quercin is insoluble in cold water, soluble in hot, of a bitter taste, and a peculiar smell. It stains paper of a yellow colour, little inferior in brilliancy to that of gamboge.

The isolation of this pigment is said to be an object of great request and importance to the dyers.

Narcotin

The Stupifying matter of opium discovered by Berdine, may be obtained by

higher
restor
Serla
meat
Ag

Haugs
Crystals
the
in lo
Ag

Liqu
not
Alter
G

emph
not to
repre

digesting opium in ether. It crystallizes in rectangular prisms with rhomboid bases. Serturner considers narcotin to be the meconate of Morphia.

Asparagin,

obtained by Robiquet and Vauquelin from the asparagus and potatoe, Crystallizes in rhomboidal prisms, stimulates the salivary glands when chewed, and dissolves in hot water.

Agaricite,

discovered by Robiquet in Liquorice, is crystalline colourless insipid insoluble in water and dissolves without alteration in Citric and Sulphuric acids.

Gentianin.

Not in the number of substances employed, and the tediousness and intricacy not to mention the absurdity of the process represented as essential to the development

of the
crash
The
the a
Sci

vision
Cap

by the
loose
publ
E

name,
pmet
delo
very p

of this pseudo-principle; I am inclined to consider it rather as an educt, than a product. The sanative efficacy of gentian resided in the alkali which I obtained from it.

Scillitin,

found by Vogel in the Squill, is bitter viscid, white, resinous and deliquescent.

Caffein,

a Crystallizable matter, observed by Robiquet in the *Coffea Arabica*, and but loosely examined by him, is white, and insoluble in Cold water.

Emelin.

A principle bearing this name, *Specacchara* is indebted for its emetic quality. Pelletier, its discoverer, described it as being white, crystalline, and very feebly if at all alkaline.

9
My
No
a Ch
hypo
and
mont
exter
the or
Lavo
ment
bade
be in
in wa
Succ
More
their

Genus IV. Oxygenia.

Proximate principles in which
oxygen is predominant.

Vegetable Acids.

It is highly culpable in
a chemist to indulge in the vanities of
hypothesis; generalization is unwarrantably
and analogy itself, unsanctioned by experi-
ment, should be employed to a very limited
extent. Thus Sheenard, by promulgating
the error in vegetable Chemistry, into which
Lavoisier had fallen, in the mineral depart-
ment: viz. that oxygen is the only acidifying
base, errs in his assumption that "if oxygen
be in a greater ratio to the hydrogen than
in water, the compound will be found acid".
Succinic, Benzoic, and ~~Propionic~~ ^{Propionic} acids contain
more hydrogen than is necessary to convert
their oxygen into water and yet they are all

Em
I do
in
ac
a.
gall
it m
impe
the
Cry
qua
prob
whod
att
h
glut
emb
the
that

eminently acid. Under this section however I shall class them, as they are analogous in their Chemical relations to the other acids.

a. Gallic acid.

This acid abounds in the gall nut, in combination with gallia from which it may be extracted by Sublimation. An aqueous infusion of galls, after a few months exposure to the atmosphere, deposits it in white filamentous Crystals. It may be obtained, in considerable quantity from the tea leaf, in which it exists probably in the state of bigallate, with an anodyne alkali, which I have frequently attempted to isolate, tho' my attempts have been as frequently abortive from the mass of glutinous extractive matter that masques and envelops it.

The most distinctive property of gallic acid is that of precipitating iron from its solutions,

and
Wyal

and C
sirop
obtain
or Sla
2. Ben

the Sy
It app
a. C

lemon
with
e. Yarl

the gro
It is L
has the

and striking with it a jet black colour.
b. Oxalic acid

is found in the opalis acedosella and Corniculata, in which it exists as a bioxalate of Potassa. It is, however, commonly obtained by reaction of Nitric acid with sugar or Starch. It crystallizes in four sided prisms.

c. Benzonic acid

is a sublimate, obtained from the Styrax benzoe and balsam of Tolu & Peru. It appears in white plumose fibres.

d. Citric acid

is ordinarily procured from lemon, or lime juice, in which it is combined with Malic acid. It is crystallizable.

e. Tartaric acid

is found in wines and in the grape in the state of bitartrate of potassa. It is susceptible of a crystalline form; and has the property of forming a pyro-tartaric

rec
of c
f. M

the
war
Bo
com
disge
that
to dis
rad,
more
with
of M

ed a f
victi
and M
City Sta

acid, which does not precipitate the acetate of lead.

f. Malic acid.

predominates in the apple, the pear, the elder, and the barberry, and in unripe fruits. It is suspected by Vogel and Bouillon Lagrange that this acid is acetic, combined with vegetable matters which disguise it — the suspicion is gratuitous. Malic acid is uncrystallizable, but affords to distillation a crystalline pyro-malic acid, and forms nearly insoluble salts with lead, Mercury, and silver — This is not the case with acetic acid.

g. Mucic acid,

Called by Scheele, Saccholactic, is a factitious acid formed by action of nitric acid with Sugar of Milk, gums, resins and Mucous. It grows in white needle-form crystals, and gives off pyro-mucic acid when

last
in Camp
we
Nitin
Y

much
soul
for

of the
Mc

religion
process
theorist
It says
Shikes
L. to be

subjected to heat.

h. Camphoric and Suberic acids
are artificially developed, by reaction of
nitric acid with camphor and cork.

i. Yungic acid,

found by Braconnot in the
mushroom combined with potash, is very
sour and deliquescent.

j. Galatrophic acid (Peltier)

Abounds in the seeds
of the Galatropa Curcas. It is uncrystallizable.

k. Meconic acid (Robiquet)

occurs in opium. The mag-
nesian precipitate, which falls during the
process for obtaining ~~opium~~ Morphia, holds
meconic acid in the state of sub-mecconate.

It crystallizes in reddish scales or plates and
strikes a red hue with ferruginous solutions.

l. Sorbic acid (Gottman)

is met with in the berries of the

So
help
of
passe
meta
leac
blan
supp
m. m.

jud
and
n. Iga

the
H
i. Me

with

Sorbus Aucuparia and in apples. It may be precipitated from the *Sorbus* by sub-acetate of lead. A current of sulphuretted hydrogen passed thro the bi-*Sorbate* of lead seizes on the metal and isolates the acid. *Sorbic acid* is scarcely crystallizable; it bears some resemblance to *Malic acid*, with which *Beacornot* supposed it to be identic.

m. *Moric* or *Moroxylic acid*

is found in an exudation from the *Morus alba*. It is white and crystalline.

n. *Igasuric acid* (Pelletier & Caventou)

Neutralizes *strychnia* in the *Mux vomica* and bears op. *I. Ipratid*.

It crystallizes in minute white needles

o. *Menispermic acid* (Boullay)

occurs in combination with *picrotoxin* in the *Cocculus Indicus*.

Kino

poppy

orn

green

callin

Kino

Succ

Prinos

Le arse

of whe

Vegeta

cytate

Acce

the Me

t. Bolle

p. Kinic acid, (Teschamps)

Observed in the extract of Quinquina, possesses a crystalline structure, and precipitates iron from its solutions, striking with it a rich green colour. By aid of heat it forms a crystalline pyrokinic acid.

q. Kinovic Acid (Blutier)

has been separated from the kina-kona.

r. Succinic acid

though generally classed among mineral acids, might with more propriety be arranged with organic products, as amber of which it is a sublimate, is evidently of vegetable origin. It forms yellow prismatic crystals of a nauseous acid taste.

s. Acetic acid

is supposed to exist in the juice of the Maple.

t. Bolitic Acid.

Beaumont had announced the

eyes
Bole
Sub
n. Tra

the a
Rup
had s
and p
with
beau
pido
romen
n. Yo

Serpe
and fo
Nicole
in so o
line in

existence of a peculiar acid principle in the
Boletus pseudoignarius. It is susceptible of
sublimation and crystallization.

u. *Krameria acid*

was found by Bischer in
the astringent root of the *Krameria triandra*.
v. *Prussic, Hydrocyanic, or Hydrazocarbureic acid*,
has been detected in bitter almonds laurel
and peach leaves, and in a number of flowers.
With ferrocyanate of potash, it furnishes a
beautiful blue colour. It is a shocking
poison, developing the most serious phe-
nomena on the nervous system.

w. To these acids I might add Quassic, Gentic,
Serpentasic, Columbic (which is crystalline
and forms a pyrocolumbic) Stropic, Laureic,
Nicotic, & Digitalic, all of which I obtained, tho'
in so small a quantity, as to preclude an exten-
sive investigation of their peculiarities.

[Faint, illegible handwriting on lined paper, likely bleed-through from the reverse side.]

P
and
hel

form
In or
part
2 gr
in wa
whic
March
Pure
heart
water
with
with
Condio

Genus V. Hydroxygenia.

Proximate principles in which oxygen and hydrogen combine in the proportion necessary to form water.

Species 1.

Fecula.

This species exists, ready formed, in such vegetables as contain it. In order to give a free exit to the Starch the parenchyma should be torn up by rasping, or grinding, and the powdered grain steeped in water and strained. This done, the liquor which passes, after ^{standing} some time, deposits the Starch.

Pure Starch is snow white, inodorous, and nearly insipid. It is not dissolved by cold water, but forms a thick gelatinoid solution with that fluid, when hot. By admixture with a caustic alkali, the solution loses its consistency, but recuperates it on the addition

of
eth
wit
Kir
Stan
Eph
a Ho
this to
Amy
Hase
Chap
from
sora
and
pro
dist
fid
is on
diff
in La

of an acid. It is insoluble in alcohol or ether, and convertible into sugar by boiling with dilute Sulphuric acid.

Kirchhoff obtained a Crystalline Sugar from Starch by long continued boiling with gluten. Exposed to a temperature of 220° , Starch acquires a blush, and exhales the odour of baked bread. To this torrefied fecula, Saussure has applied the term Amydine; and Caumont maintains that Starch gelatinized with hot water undergoes a similar change and is identic with amydine: 1^o because from neither of these States can Starch be recovered; 2^o from a presumption that it is amydine and not Starch which has the property of producing a blue colour with iodine. This distinguished Chemist does not speak from experiment. His first statement is too general and vague; his second is erroneous — for if a small portion of Iodine be diffused thro cold water, in a glass vessel holding Starch in suspension, and left quiescent for some time,

[Faint, illegible handwriting on lined paper, likely bleed-through from the reverse side.]

it w
of co
1472

last
1472
Tha

from
ana
by
Pro

Law
was
a b
The

Orin

it will acquire an elegant blue colour, the result of iodine action on Starch.

Hordein,

A variety of Glucan, observed by Roust in barley. It is a yellow acid powder, convertible into Starch and Sugar by the germination of barley.

Tragopyrein.

This species of Starch I obtained from Buckwheat (*Tragopyrum*). It is somewhat analogous to Hordein and may be distinguished by its solubility in hot alcohol.

Oryzein,

An amylaceous substance which I ~~xxx~~ procured from rice, diffuses in cold water without forming a gelatine and strikes a blue with iodine.

Zein,

A peculiar Starch existing in Indian corn, is soluble in proof spirit.

Wm

is so

in wa

tilwa

etwa

is the

Inu

leca

gadu

ation

and is

lup

y

insol

is a

trucid

Ulm, in

found in an exudation from the elm,
is solid, tasteless, black, and brilliant, soluble
in water and insoluble in alcohol. Pure potassa
trituated with saw dust furnishes ulmin, by
abstracting from lignin oxygen and hydrogen
in the proportion of water.

Malin,

extracted by Boe from the root of the
Elecampane, dissolves in hot water without
producing viscosity, and precipitates on refrige-
ration. It strikes a green colour with iodine
and is dissolved without decomposition by
Sulphuric acid.

Species 2

Gums,

in a state of purity, are
insoluble in alcohol and ether, soluble in the
oils and in water forming with the latter
mucilage, and passing speedily from their

[Faint, mirrored handwriting, likely bleed-through from the reverse side of the page. The text is illegible due to fading and mirroring.]

aque
By
acid,
reac
how
of the
Cera

d are
in the
Sa

rema
and
but so

the
at com

aqueous solution into acetous fermentation.
By distillation they furnish an impure acetic acid, which has been called pyromucic, and by reaction with nitric acid their elements form new combinations, attended by the production of Malic, Mucic, and oxalic acids.

Cerasin,

A generic term applied to such gums as are insoluble in cold water, but analogous in every other respect to common gum.

Sacogummite,

A sweet gummoid matter remarked by Robiquet in liquorice. It is yellow and solid insoluble in cold water and alcohol but soluble in both these fluids when hot.

Species 3.

Sugar.

The principle which forms the nutritious portion of the vegetable secretions, is commonly extracted from the arundo

Sal
exid
and
Calle
lath
Whe
gela
Cryp
or in
Solu
Solu
by pr
Wale
Sug
of re
oped
of Cop
redul
hence
ra as

Saccharifera or sugar cane in which there exist two distinct varieties viz; a crystallizable, and an uncrystallizable sugar. The former is called raw sugar, Muscovado, loaf sugar^{or}; the latter Melasses or treacle.

When sugar is perfectly detached from treacle, gelatine, gluten, and saccho-mucous matter, it crystallizes in incomplete coniciform octoëdra, or in tetrahedral prisms. It is diaphanous, soluble in all proportions in water, slightly soluble in alcohol and becomes phosphorescent by friction. Long exposure to the heat of boiling water, robs it of the property of crystallizing.

Sugar, dissolved in hot water, has the property of restoring to the metallic state several oxidized metals, and metallic salts. The oxides of Copper, so pernicious to animal life, are reduced and rendered innocent by sugar—hence it may be advantageously exhibited as an antiloimic.

Ma

of de
act

It is

but

the

some

and

of

Aug

max

or h

but

may

to

or

phy

Mannite.

Sugar of Manna has the property of dissolving the yellow oxide of lead, and of acting like an alkali on vegetable blues. It is soluble in boiling water and alcohol, but on cooling falls in crystalline flocks.

Gungite.

Sugar of Mushrooms possesses some peculiarities. It is white crystalline and efflorescent.

Honey

is by some considered a vegetable sugar, by others an animal product. That it undergoes some modification in the stomach or honey bag of the bee, scarce admits of a doubt; but that it is essentially of vegetable origin may be demonstrated, as well by the untutored peasant who sucks it from the clover blossom or observes it in the honey dew, as by the physiologist who demonstrates on the plant

the
who
has
inde
Glor
plac
hum
wher
flore
Rom
lexic
honey
occas
Main
Hone
and
sugar
acid
La

the nectary, the melliferous glands, and the whole complicated apparatus which nature has constructed for the defence of a fluid so indispensable to the vegetable economy. Honey, obtained from poisonous or narcotic plants, retains their deleterious qualities. Numerous examples of its toxic agency, when procured from hemlock "gelide collection flore Cicutæ", are recorded by the Greek and Roman writers. It is stated by Porro, a lexicographer of the 15th Century, that the honey of bees frequenting the Rhododendron occasions a peculiar kind of Madness, called *Manimemon*.

Honey, like sugar, consists of a crystallizable and an uncrystallizable portion; and, like sugar is convertible into malic and oxalic acids, by reaction with nitric acid.

Sarcocollin,

Obtained from the *Sarcocoll*

[Faint, illegible handwriting on lined paper, likely bleed-through from the reverse side.]

it o
fillor
flin

line,
lars


skell
to th
alrea
suffe
the g
it is
it is

is brown, brittle, unctuous, and of a sweetish
bitter taste.

Fluide,

observed by Pelletier in the gum-resin of the
olive, is white, brilliant, and Crystalloid, tasting like
Saccharin.

Thus have I completed a rapid and succinct
Sketch of a Science, which owes its existence
to the nineteenth century; one which has
already rendered the most signal services to
suffering humanity; and one, which holds out
the golden promise of becoming the most important
as it unquestionably is the most useful, branch
of its generic Science.



See

Re.

5